

Supplement zum Buchbeitrag

„Evidenzbasierte Gesundheitsinformation“

Autoren: Albrecht M, Mühlhauser I, Steckelberg A

Literaturangaben zum Kapitel V 1.2

1.2.1 Darstellung von Zahlen

- Akl, E. A., Oxman, A. D., Herrin, J., Vist, G. E., Terrenato, I., Sperati, F., Costiniuk, C., Blank, D. & Schünemann, H. (2011). Using alternative statistical formats for presenting risks and risk reductions. *Cochrane Database Syst Rev*, Issue 3, No.: CD006776. doi:10.1002/14651858.CD006776.pub2
- Berry, D. C., Knapp, P. & Raynor, T. (2006). Expressing medicine side effects: assessing the effectiveness of absolute risk, relative risk, and number needed to harm, and the provision of baseline risk information. *Patient Education and Counseling*, 63, 89-96. doi:10.1016/j.pec.2005.09.003
- Carling, C. L., Kristoffersen, D. T., Montori, V. M., Herrin, J., Schünemann, H. J., Treweek, S., Akl, E. A. & Oxman, A. D. (2009). The effect of alternative summary statistics for communicating risk reduction on decisions about taking statins: a randomized trial. *PLoS Medicine* 6, e1000134. doi:10.1371/journal.pmed.1000134
- Knapp, P., Gardner, P. H., Carrigan, N., Raynor, D. K. & Woolf, E. (2009). Perceived risk of medicine side effects in users of a patient information website: a study of the use of verbal descriptors, percentages and natural frequencies. *British Journal of Health Psychology*, 14, 579-594. doi:10.1348/135910708X375344
- Natter, H. M. & Berry, D. C. (2005). Effects of presenting the baseline risk when communicating absolute and relative risk reductions. *Psychology, Health & Medicine*, 10, 326-334. doi:10.1080/13548500500093407
- Ruiz, J. G., Andrade, A. D., Garcia-Retamero, R., Anam, R., Rodriguez, R. & Sharit, J. (2013). Communicating global cardiovascular risk: Are icon arrays better than numerical estimates in improving understanding, recall and perception of risk? *Patient Education and Counseling*, 93, 394-402. doi: 10.1016/j.pec.2013.06.026
- Schwartz, L. M., Woloshin, S., Black, W. C. & Welch, H. G. (1997). The role of numeracy in understanding the benefit of screening mammography. *Annals of Internal Medicine*, 127(11):966-972. doi:10.7326/0003-4819-127-11-199712010-00003
- Sheridan, S. L., Pignone, M. P. & Lewis, C. L. (2003). A randomized comparison of patients' understanding of number needed to treat and other common risk reduction formats. *Journal of General Internal Medicine*, 18, 884-892. doi: 10.1046/j.1525-1497.2003.21102.x

Woloshin, S. & Schwartz, L. M. (2011). Communicating data about the benefits and harms of treatment: a randomized trial. *Annals of Internal Medicine*, 155, 87-96. doi:10.7326/0003-4819-155-2-201107190-00004

1.2.2 Verbale Darstellung versus Zahlen

Berry, D. C., Knapp, P. & Raynor, D. K. (2002). Provision of information about drug side effects to patients. *Lancet*, 359, 853-854. doi:10.1016/S0140-6736(02)07923-0

Berry, D. C., Knapp, P. & Raynor, T. (2002). Is 15 per cent very common? Informing people about the risks of medication side effects. *International Journal of Pharmacy Practice*, 10, 145-151. doi:10.1111/j.2042-7174.2002.tb00602.x

Berry, D. C., Raynor, T., Knapp, P. & Bersellini, E. (2004). Over the counter medicines and the need for immediate action: a further evaluation of European Commission recommended wordings for communicating risk. *Patient Education and Counseling*, 53, 129-134. doi:10.1016/S0738-3991(03)00111-3

Berry, D. C., Raynor, D. K. & Knapp, P. (2003). Communication risk of medication side effects: an empirical evaluation of EU recommended terminology. *Psychology, Health & Medicine*, 8, 251-263. doi:10.1080/1354850031000135704

Bundesinstitut für Arzneimittel und Medizinprodukte (2013). Wie sollen die Häufigkeiten für Nebenwirkungen in der Produktinformation angegeben werden? Zugriff unter: http://www.bfarm.de/SharedDocs/4_FAQ/DE/Arzneimittel/pal/ja-ampal-faq.html?nn=1010882

European Commission (1998): A guideline on the readability of the label and package leaflet of medicinal products for human use. European Commission, Directorate-General III.

Knapp, P., Gardner, P. H., Carrigan, N., Raynor, D. K. & Woolf, E. (2009). Perceived risk of medicine side effects in users of a patient information website: a study of the use of verbal descriptors, percentages and natural frequencies. *British Journal of Health Psychology*, 14, 579-594. doi:10.1348/135910708X375344

Knapp, P., Raynor, D. K. & Berry, D. C. (2004). Comparison of two methods of presenting risk information to patients about the side effects of medicines. *Quality & Safety in Health Care*, 13, 176-180. doi:10.1136/qshc.2003.009076

Knapp, P., Raynor, D. K., Woolf, E., Gardner, P. H., Carrigan, N. & McMillan, B. (2009). Communication the risk of side effects to patients. An evaluation of UK regulatory recommendations. *Drug Safety*, 32, 837-849. doi:10.2165/11316570-000000000-00000

Man-Son-Hing, M., O'Connor, A. M., Drake, E., Biggs, J., Hum, V. & Laupacis, A. (2002). The effect of qualitative vs. quantitative presentation of probability estimates on patient decision-making: a randomized trial. *Health Expectations*, 5, 246-255. doi:10.1046/j.1369-6513.2002.00188.x

Marteau, T. M., Saidi, G., Goodburn, S., Lawton, J., Michie, S. & Bobrow, M. (2000). Numbers or words? A randomized controlled trial of presenting screen negative results to pregnant women. *Prenatal Diagnosis*, 20 (9), 714-718.

Ziegler, A., Hadlak, A., Mehlbeer, S. & König, I. R. (2013). Verständlichkeit von Nebenwirkungsrisiken im Beipackzettel. Eine Umfrage unter Ärzten, Apothekern und Juristen. *Deutsches Ärzteblatt International*, 110, 669-673. doi:10.3238/arztebl.2013.0669

1.2.3 Grafische Darstellungen

Ancker, J. S., Senathirajah, Y., Kukafka, R. & Starren, J. B. (2006). Design features of graphs in health risk communication: a systematic review. *Journal of the American Medical Informatics Association*, 13, 608-618. doi:10.1197/jamia.M2115

Brewer, N. T., Gilkey, M. B., Lillie, S. E., Hesse, B. W. & Sheridan, S. L. (2012). Tables or bar graphs? Presenting test results in electronic medical records. *Medical Decision Making*, 32, 545-453. doi:10.1177/0272989X12441395

Bunge, M., Mühlhauser, I. & Steckelberg, A. (2010). What constitutes evidence-based patient information? Overview of discussed criteria. *Patient Education and Counseling*, 78, 316-328. doi:10.1016/j.pec.2009.10.029

Feldman-Stewart, D., Brundage, M. D. & Zotov, V. (2007). Further insight into the perception of quantitative information: judgments of gist in treatment decisions. *Medical Decision Making*, 27, 34-43. doi:10.1177/0272989X06297101

Ghosh, K., Crawford, B. J., Pruthi, S., Williams, C. I., Neal, L., Sandhu, N. P., Johnson, R. E., Wahner-Roedler, D., Britain, M. K., Cha, S. S. & Ghosh, A. K. (2008). Frequency format diagram and probability chart for breast cancer risk communication: a prospective, randomized trial. *BMC Womens Health*, 8, 18. doi:10.1186/1472-6874-8-18

Hawley, St., Zikmund-Fisher, B., Ubel, P., Jancovic, A., Lucas, T. & Fagerlin, A. (2008). The impact of the format of graphical presentation on health-related knowledge and treatment choices. *Patient Education and Counseling*, 73, 448-455.

Heesen, C., Köpke, S., Kasper, J., Richter, T., Beier, M. & Mühlhauser I. (2013). Immuntherapien der Multiplen Sklerose. Zugriff unter: http://www.ms-netz-hamburg.de/fileadmin/msnetzhamburg/dokumente/ISDIMS_Neufassung_Maerz_2013.pdf

Lee, D. H. & Mehta, M. D. (2003). Evaluation of a visual risk communication tool: effects on knowledge and perception of blood transfusion risk. *Transfusion*, 43, 779-787.

Ruiz, J. G., Andrade, A. D., Garcia-Retamero, R., Anam, R., Rodriguez, R. & Sharit, J. (2013). Communicating global cardiovascular risk: Are icon arrays better than numerical estimates in improving understanding, recall and perception of risk? *Patient Education and Counseling*, 93, 394-402.

Sprague, D., LaVallie, D. L., Wolf, F. M., Jacobsen, C., Sayson, K. & Buchwald, D. (2011). Influence of graphic format on comprehension of risk information among American Indians. *Medical Decision Making*, 31, 437. doi:10.1177/0272989X10391096.

Steckelberg, A. & Mühlhauser, I. (2011): Darmkrebs – Früherkennung. Zugriff unter: <http://www.gesundheit.uni-hamburg.de/upload/NeueDarmkrebsbroschuere2011.pdf>

- Tait, A. R., Voepel-Lewis, T., Brennan-Martinez, C., McGonegal, M. & Levine, R. (2012). Using animated computer-generated text and graphics to depict the risks and benefits of medical treatment. *American Journal of Medicine*, 125, 1103-1110.
- Tait, A. R., Voepel-Lewis, T., Zikmund-Fisher, B. J. & Fagerlin, A. (2010): Presenting research risks and benefits to parents: Does format matter? *Anesthesia and Analgesia*, 111, 718-723.
- Tait, A. R., Voepel-Lewis, T., Zikmund-Fisher, B. J. & Fagerlin, A. (2010). The effect of format on parents' understanding of the risks and benefits of clinical research: a comparison between text, tables, and graphics. *Journal of Health Communication*, 15, 487-501.
- Zikmund-Fisher, B. J., Ubel, P. A., Smith, D. M., Derry, H. A., McClure, J. B., Stark, A., Pitsch, R. K. & Fagerlin, A. (2008). Communicating side effect risks in a tamoxifen prophylaxis decision aid: the debiasing influence of pictographs. *Patient Education and Counseling*, 73, 209-214.

1.2.4 Abbildungen und Piktogramme

- Boelhouwer, E., Davis, J., Franco-Watkins, A., Dorris, N. & Lungu, C. (2013). Comprehension of hazard communication: effects of pictograms on safety data sheets and labels. *Journal of Safety Research*, 46, 145-155.
- Bunge, M., Mühlhauser, I. & Steckelberg, A. (2010). What constitutes evidence-based patient information? Overview of discussed criteria. *Patient Education and Counselling*, 78, 316-328. doi:10.1016/j.pec.2009.10.029
- King, S. R., Mc Caffrey, D. J., Bentley, J. P. Bouldin, A., Hallam, J. & Wilkin, N. E. (2012). The influence of symbols on the short-term recall of pharmacy generated prescription medication information in a low health literate sample. *Journal of Health Communication*, 17, 280-293.
- Mansoor, L. E. & Dowse, R. (2003). Effect of pictograms on readability of patient information materials. *Annals of Pharmacotherapy*, 37, 1003-1009.
- Roundtable on Health Literacy (2008). Board on Population Health and Public Health Practice, Institute of Medicine. In L. M. Hernandez (ed.), Standardizing medication labels: confusing patients less. Workshop summary. Washington DC: National Academies Press.
- Sahm, L. J., Wolf, M. S., Curtis, L. M. Behan, R., Brennan, M., Gallwey, H., Mc Carthy, S. (2012). What's in a label? An exploratory study of patient-centered drug instructions. *European Journal of Clinical Pharmacology*, 68, 777-782.
- Thompson, A. E., Goldszmidt, M. A. & Schwartz, A. J. (2010). A randomized trial of pictorial versus prose-based medication information pamphlets. *Patient Education and Counseling*, 78, 389-393.
- Yin, H. S., Mendelsohn, A. L. & Fiermann, A. (2011). Use of a pictographic diagram to decrease parent dosing errors with infant acetaminophen: a health literacy perspective. *Academic Pediatrics*, 11, 50-57.

1.2.5 Persönliche Erfahrungsberichte (Narrative)

- Bandura, A. (2004). Health promotion by social cognitive means. *Health Education & Behavior*, 31, 143-164.
- Betsch, C., Renkewitz, F. & Haase, N. (2012). Effect of narrative reports about vaccine adverse events and bias-awareness disclaimers on vaccine decisions: a simulation of an online patient social network. *Medical Decision Making*, 33, 14-25.
- Betsch, C., Ulshöfer, C., Renkewitz, F. & Betsch, T. (2011): The influence of narrative vs. statistical information on perceiving vaccination risks. *Medical Decision Making*, 31, 742-753. doi:10.1177/0272989X11400419
- Bollinger, S. & Kreuter, M. W. (2012). Real-time moment-to-moment emotional responses to narrative and informational breast cancer videos in African American women. *Health Education Research*, 27, 537-543.
- Bunge, M., Mühlhauser, I. & Steckelberg, A. (2010). What constitutes evidence-based patient information? Overview of discussed criteria. *Patient Education and Counselling*, 78, 316-328. doi:10.1016/j.pec.2009.10.029
- Cody, R. & Lee, C. (1990). Behaviors, beliefs, and intentions in skin cancer prevention. *Journal of Behavioral Medicine*, 13, 373-388.
- De Wit, J. B., Das, E. & Vet, R. (2008). What works best: objective statistics or a personal testimonial? An assessment of the persuasive effects of different types of message evidence on risk perception. *Health Psychology*, 27, 110-115.
- Dillard, A.J., Fagerlin, A., Dal Cin, S., Zikmund-Fisher, B.J. & Ubel, P. A. (2010). Narratives that address affective forecasting errors reduce perceived barriers to colorectal cancer screening. *Social Science & Medicine*, 71, 45-52. doi:10.1016/j.socscimed.2010.02.038
- Elwyn, G., O'Connor, A., Stacey, D., Volk, R., Edwards, A., Coulter, A., Thomson, R., Barratt, A., Barry, M., Bernstein, S., Butow, P., Clarke, A., Entwistle, V., Feldman-Stewart, D., Holmes-Rovner, M., Llewellyn-Thomas, H., Moumjid, N., Mulley, A., Ruland, C., Sepucha, K., Sykes, A., Whelan, T. & International Patient Decision Aids Standards (IPDAS) Collaboration. (2006). International patient decision aids standards (IPDAS) collaboration. Developing a quality criteria framework for patient decision aids: online international delphi consensus process. *BMJ*, 333, 417. doi: 10.1136/bmj.38926.629329.AE
- Fagerlin, A., Wang, C. & Ubel, P. (2005). Reducing the influence of anecdotal reasoning on people's health care decisions: is a picture worth a thousand statistics? *Medical Decision Making*, 25, 398-405.
- Feldman-Stewart, D., Brennenstuhl, S., McIsaac, K., Austoker, J., Charvet, A., Hewitson, P., Sepucha, K. R. & Whelan, T. (2007). A systematic review of information in decision aids. *Health Expectations*, 10, 46-61. doi: 10.1111/j.1369-7625.2006.00420.x
- Greene, K. & Brinn, L. S. (2003). Messages influencing college women's tanning bed use: statistical versus narrative evidence format and a self-assessment to increase perceived susceptibility. *Journal of Health Communication*, 8, 443-461.

- Khangura, S., Bennett, C., Stacey, D. & O'Connor A. M. (2008). Personal stories in publicly available patient decision aids. *Patient Education and Counseling*, 73, 456-464.
doi:10.1016/j.pec.2008.07.035
- Kreuter, M. W., Holmes, K., Alcaraz, K., Kalesan, B., Rath, S., Richert, M., McQueen, A., Caito, N., Robinson, L. & Clark, E. M. (2010). Comparing narrative and informational videos to increase mammography in low-income African American women. *Patient Education and Counseling*, 81S, S6-S14.
- Mazor, K. M., Baril, J., Dugan, E., Spencer, F., Burgwinkle, P. & Gurwitz, J. H. (2007). Patient education about anticoagulant medication: is narrative evidence or statistical evidence more effective? *Patient Education and Counseling*, 69, 145-157.
- McDonald, D. D., Goncalves, P. H., Almarino, V. E., Krajewski, A. L., Cervera, P. L., Kaeser, D. M., Lillvik, C. A., Sajkowicz, T. L. & Moose, P. E. (2006). Assisting women to learn myocardial infarction symptoms. *Public Health Nursing*, 23, 216-223.
- Prati, G., Pietrantoni, L. & Zani, B. (2012). Influenza vaccination: the persuasive-ness of messages among people aged 65 years and older. *Health Communication*, 27, 413-420.
- Ricketts, M., Shanteau, J., McSpadden, B. & Fernandez-Medina, K. M. (2010). Using stories to battle unintentional injuries: narratives in safety and health communication. *Social Science & Medicine*, 70, 1441-1449.
- Rook, K. S. (1987). Effects of case history versus abstract information on health attitudes and behaviors. *Journal of Applied Social Psychology*, 17, 533-553.
- Rook, K. S. (1986). Encouraging preventive behavior for distant and proximal health threats: effects of vivid versus abstract information. *Journal of Gerontology*, 41, 526-534.
- Shaffer, V. A. & Zikmund-Fisher, B. J. (2012). All stories are not alike: a purpose-, content-, and valence-based taxonomy of patient narratives in decision aids. *Medical Decision Making*, 33, 4-13. doi:10.1177/0272989X12463266
- Slater, M. D., Buller, D. B., Waters, E., Archibeque, M. & LeBlanc, M. (2003). A test of conversational and testimonial messages versus didactic presentations of nutrition information. *Journal of Nutrition Education and Behavior*, 35, 255-259.
- Ubel, P. A., Jepson, C. & Baron, J. (2001). The inclusion of patient testimonials in decision aids: effects on treatment choices. *Medical Decision Making*, 21, 60-68.
- Winterbottom, A., Bekker, H. L., Conner, M. & Mooney, A. (2008). Does narrative information bias individual's decision making? A systematic review. *Social Science & Medicine*, 67, 2079-2088.
doi:10.1016/j.socscimed.2008.09.037

1.2.6 Zielgruppenspezifische Sprache

- Bates, B. R., Romina, S. M. & Ahmed, R. (2007): The effect of improved readability scores on consumers' perceptions of the quality of health information on the internet. *Journal of Cancer Education*, 22, 15-20.
- Berry, D. C., Michas, I. C. & Bersellini, E. (2003). Communicating information about medication: the benefits of making it personal. *Psychology & Health*, 18, 127-139.
- Dowe, M. C., Lawrence, P. A., John Carlson, J. & Keyserling, T. C. (1997). Patients' use of health-teaching materials at three readability levels. *Applied Nursing Research*, 10, 86-93.
- Europäische Union (2013). Klar und deutlich schreiben (deutsche Fassung des Leitfadens). Zugriff unter: http://ec.europa.eu/translation/german/german_en.htm
- Hoffmann, T. & Worrall, L. (2004). Designing effective written health education materials: considerations for health professionals. *Disability and Rehabilitation*, 26, 1166-1173. doi:10.1080/09638280410001724816
- Meade, C. D., Byrd, J. C. & Lee, M. (1989). Improving patient comprehension of literature on smoking. *American Journal of Public Health*, 79, 1411-142.
- Perera, K. Y., Ranasinghe, P., Adikari, A. M., Balagobi, B., Constantine, G. R., Jayasinghe, S. (2012). Medium of language in discharge summaries: Would the use of native language improve patients' knowledge of their illness and medications? *Journal of Health Communication*, 17, 141-148.
- Sansgiry, S. S., Chanda, S. & Shringarpure, G. S. (2007). Impact of bilingual product information labels on Spanish-speaking adults' ability to comprehend OTC information. *Research in Social & Administrative Pharmacy*, 3, 410-425.
- Stableford, S. & Mettger, W. (2007). Plain language: a strategic response to the health literacy challenge. *Journal of Public Health Policy*, 28, 71-93. doi:10.1057/palgrave.jphp.3200102
- Zite, N. B. & Wallace, L. S. (2011). Use of a low-literacy informed consent form to improve women's understanding of tubal sterilization. A randomized controlled trial. *Obstetrics and Gynecology*, 117, 1160-1166.